



# Axill Solutions Glufosinate 280SL

Axill Solutions Glufosinate 280SL is a non-selective herbicide. Apply Axill Solutions Glufosinate 280SL for post emergence control of listed weed species in listed berry, tree, and vine crops. Axill Solutions Glufosinate 280SL may also be applied for potato vine desiccation. Axill Solutions Glufosinate 280SL is also a non-selective herbicide for hooded spray applications in conventional cotton and post emergence broadcast use on canola, field corn, sweet corn\*, cotton, soybean, and sugar beets\* designated as glufosinate-resistant. Axill Solutions Glufosinate 280SL may be applied as a broadcast burndown application before planting or prior to emergence of any conventional or glufosinate-resistant variety of canola, field corn, cotton, soybean, or sugar beet. Axill Solutions Glufosinate 280SL may also be applied for canola, corn, cotton, and soybean seed propagation.

\*Not for use in California

<b>Active Ingredient:</b>	<b>% By Weight</b>
Glufosinate-ammonium .....	24.5%
<b>Other Ingredients:</b> .....	75.5%
<b>Total:</b> .....	<b>100.0%</b>

Contains 2.34 lbs. of active ingredient per gallon.

## KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you **DO NOT** understand this label, find someone to explain it to you in detail.)

### FIRST AID

#### IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

#### IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

#### IF SWALLOWED:

- Immediately call a poison control center or doctor.
- **DO NOT** induce vomiting unless told to do so by a poison control center or doctor.
- Have person sip a glass of water if able to swallow.
- **DO NOT** give anything by mouth to an unconscious person.

### NOTE TO PHYSICIAN

If this product is ingested, endotracheal intubation and gastric lavage must be performed as soon as possible, followed by charcoal and sodium sulfate administration.

### HOTLINE NUMBERS

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For 24-Hour Medical Emergency Assistance (Human or Animal), call: **1-800-222-1222**. For Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), call CHEMTREC: **1-800-424-9300**. For general information on this product contact the National Pesticide Information Center (NPIC) at 1-800-858-7378, Monday-Friday, 8am – 12pm PST, or at <http://npic.orst.edu>.

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

EPA Reg. No.: 93809-14

EPA Est. No.: 83411-MN-001

**Manufactured For:**  
Axill Solutions, LLC  
422 Jasmine Way  
Roseburg, OR 97471

**Net Contents: 265 Gallons**

# PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### WARNING

Causes substantial but temporary eye injury. Harmful if absorbed through skin. Harmful if swallowed. **DO NOT** get in eyes. Avoid contact with skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves including barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, polyvinyl chloride (PVC)  $\geq 14$  mils, or Viton®  $\geq 14$  mils
- Shoes plus socks
- Protective eyewear (goggles, face shield or safety glasses)
- Wear a chemical-resistant apron when mixing/loading and cleaning equipment.

#### Applicators using ground boom equipment with open cabs to treat cotton must wear:

- Long-sleeved shirts and long pants
- Shoes and socks
- Chemical-resistant gloves

#### Mixer/loaders supporting ground boom applications to corn, canola, soybean, cotton, citrus fruit, pome fruit, stone fruit, and olives must wear:

- Long-sleeved shirts and long pants
- Shoes and socks
- Chemical-resistant gloves

#### Mixers/loaders supporting aerial applications must wear:

- A minimum of a NIOSH approved filtering face piece respirator with any N filter (TC-84A). You can also use other NIOSH approved particulate respirators that offer more protection.
- When mixing and loading wear a chemical-resistant apron.
- For overhead exposure wear chemical-resistant headgear.
- When cleaning equipment wear a chemical-resistant apron.

### ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY REQUIREMENTS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

**DO NOT** apply directly to water or to areas where surface water is present. **DO NOT** apply to intertidal areas below the mean highwater mark. **DO NOT** contaminate water by cleaning of equipment or disposal of equipment wash waters or rinsate.

This pesticide is toxic to vascular plants. Use in strict accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils with shallow groundwater. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several weeks after application. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of glufosinate-ammonium from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

### PHYSICAL-CHEMICAL HAZARDS

**DO NOT** mix or allow coming in contact with oxidizing agents or hazardous chemical reaction will occur.

# DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** use this product until you have read the entire label. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

In the State of New York Only: Not For Use In Nassau and Suffolk Counties.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses; and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry-interval (REI) of 12 hours, with the following exceptions:

- The REI for works engaged in scouting activities in corn, canola, and soybeans is 4 days.
- The REI for workers to move irrigation piping is 7 days for all crops.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants
- Chemical-resistant gloves including barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, polyvinyl chloride (PVC)  $\geq 14$  mils, or Viton<sup>®</sup>  $\geq 14$  mils
- Shoes plus socks
- Protective eyewear (goggles, face shield or safety glasses)

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The application for NON-CROP USES are not within the scope of the WPS.

Keep unprotected persons out of treated areas until sprays have dried.

## PRODUCT INFORMATION

**Axill Solutions Glufosinate 280SL** is a water soluble herbicide for use as a foliar spray in bushberries (crop subgroup 13B), trees: citrus (crop group 10-10), olive, pome fruit (crop group 11-10), stone fruit (crop group 12-12), tree nut (crop group 14-12), vines (grapes); also for foliar spray in glufosinate-resistant canola, corn\*, cotton, soybean and sugar beet\*, as well as hooded spray in non-glufosinate-resistant cotton; and for use in potato desiccation, burndown, non-crop and fallow fields. **Axill Solutions Glufosinate 280SL** exhibits foliar activity to a wide spectrum of broadleaf, grass, annual and perennial weeds.

\*Not for use on sweet corn or sugar beet in California.

When making application to bushberries (crop subgroup 13B), trees: citrus (crop group 10-10), olive, pome fruit (crop group 11-10), stone fruit (crop group 12-12), tree nut (crop group 14-12), vines (grapes), keep **Axill Solutions Glufosinate 280SL** solution from contacting green bark, branches or vegetation, to prevent injury to plants. Trunks with callused, established brown bark, or shielded by nonporous wraps, grow tubes or waxed containers can be sprayed with **Axill Solutions Glufosinate 280SL**.

When making post-emergent foliar application to canola, corn, cotton, soybean and sugar beet, be sure to make application only to glufosinate-resistant crops. If used on row crops not designated as glufosinate-resistant, Axill Solutions, LLC does not warrant that they will tolerate the use of this product without harm to the extent consistent with applicable law.

Glufosinate-resistant crops contain a gene which results in a plant that is resistant to glufosinate-ammonium, whereas other crops **DO NOT** contain this gene and will suffer serious crop injury or death. When applying **Axill Solutions Glufosinate 280SL**, take care that spray does not contact desirable vegetation, foliage, or green tissue of non-glufosinate-resistant plants.

When making application to conventional cotton, use a hooded sprayer, to avoid exposure of cotton plants to **Axill Solutions Glufosinate 280SL**. If **Axill Solutions Glufosinate 280SL** comes in contact with non-glufosinate-resistant cotton plants (foliage or stems), serious injury or loss of plant could occur.

**Axill Solutions Glufosinate 280SL** can be applied broadcast prior to planting or emergence of conventional or glufosinate-resistant canola, corn, cotton, soybean or sugar beet, to act as a burndown agent for existing weed species.

## IMPORTANT

- **Axill Solutions Glufosinate 280SL** has foliar activity, but little to no activity in soil. Apply product to actively growing weeds for best control. Little to no control of weeds that emerge after application will be achieved.
- For optimum weed control:
  - Make sure uniform, thorough spray coverage is obtained.
  - Avoid cultivation from 5 days prior to application to 7 days following application.
  - If possible, avoid application in heavy dew, fog, mist, or rain.
  - Apply **Axill Solutions Glufosinate 280SL** between dawn and 2 hours before sunset (to optimize lambsquarters and velvetleaf control).
- When applying to most weed species, **Axill Solutions Glufosinate 280SL** will be rainfast 4 hours after treatment (if weeds are exposed to rain prior to 4 hours after application, may need retreatment, or may give decreased weed control)
- Decreased weed control may be observed if weeds are under stress due to such environmental conditions as cloudy weather, cool temperatures, or drought.

## MIXING DIRECTIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Before adding **Axill Solutions Glufosinate 280SL** to the spray tank, make sure the tank has been thoroughly cleaned and free of any product residue. See the **CLEANING** section for instructions. The spray tank must also be properly calibrated prior to use.

**Axill Solutions Glufosinate 280SL** dilutes quickly and easily in water. For proper mixing fill the spray tank with water and begin agitation. Add **Axill Solutions Glufosinate 280SL** and continue agitation. Prior to applying **Axill Solutions Glufosinate 280SL** to crops or use sites, flush out the spray system lines. This will guarantee that the complete system contains and is applying the correct concentration of **Axill Solutions Glufosinate 280SL**.

### Tank Mixtures – Compatibility

If applying **Axill Solutions Glufosinate 280SL** in a tank mix or with other mixing products, test compatibility prior to mixing in application equipment.

**Compatibility Test:** In a lidded jar (~1 quart size), add all mix partners, in their relative proportions. Invert, shake or mix the jar thoroughly. If mixture forms precipitates (flakes or sludge), gels, balls up or forms oily film or layers, this indicates incompatibility. Let the mixture stand for 15 minutes before determining compatibility.

### Tank Mixtures – Mixing

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Axill Solutions Glufosinate 280SL** can be applied on its own or in combination with other herbicides to provide additional residual control or to support product performance. Make certain that any tank mix partners are registered for use on the same crops or use sites on this product label, and follow all label instructions, restrictions, and precautions on both labels.

Consult specific crop use directions for additional information regarding rates and restrictions.

Mix **Axill Solutions Glufosinate 280SL** using the following procedure:

1. Fill a clean spray tank with half of the water required for treatment.
2. Begin agitation.
3. If mixing with a flowable/wettable powder tank mix partner, use a clean container to make a slurry of the powder and a small amount of water. Add the slurry to the spray tank; rinse slurry mix container; add rinsate to spray tank.
4. If required, add appropriate amount of ammonium sulfate (AMS) to the spray tank.
5. If mixing with a liquid tank mix partner, add to spray tank next.
6. Add the rest of the water to the spray tank.
7. Continue agitation and add measured amount of **Axill Solutions Glufosinate 280SL** to spray tank.
8. A silicone antifoam agent can be added, if needed to deter foaming.

Maintain agitation until the product is used. If spray mixture settles, be sure to thoroughly agitate to remix the solution prior to continuing application. Foaming can be minimized by positioning bypass line near or on the bottom of the tank. Use screen size of 50 mesh or greater in line strainers and nozzles.

## APPLICATION DIRECTIONS

Apply **Axill Solutions Glufosinate 280SL** to weeds in a manner that results in good coverage of the weed foliage, otherwise diminished weed control will result. Make application when weeds are small and growing (3 inches or less in height), at rates indicated in crop specific sections of this product label.

## Ground Application

**Axill Solutions Glufosinate 280SL** can be applied as a broadcast treatment. To ensure complete spray coverage, mix product in at least 15 gallons of water per acre, and up to 20 gallons of water per acre if weed or crop canopy is heavy, unless otherwise indicated in specific crop use directions. For additional information concerning product application, consult the **MANDATORY SPRAY DRIFT MANAGEMENT** portion of this label.

## Aerial Application

Unless otherwise specified in specific crop use directions, **Axill Solutions Glufosinate 280SL** can be applied aerially. Mix product in a minimum of 10 gallons of water per acre. For additional information concerning product application, consult the **MANDATORY SPRAY DRIFT MANAGEMENT** portion of this label.

**Application Restriction:** Air-assisted spray equipment, flood jet nozzles or controlled droplet application equipment cannot be used.

## CLEANING

Before storing, mixing or applying **Axill Solutions Glufosinate 280SL**, ensure that all tanks or storage containers have been thoroughly cleaned, including all lines and filters. Thoroughly clean and rinse any containers with a commercial tank cleaner, and consult labeling for any product previously contained or used for additional specific cleaning directions.

After storing, mixing or applying **Axill Solutions Glufosinate 280SL**, clean the equipment or containers thoroughly:

- Triple rinse spray equipment
- Use a commercial tank cleaner following rinses
- Completely remove any foam or rinsate from the boom and spray tank
- Consult pesticide disposal directions for disposing of rinsate

## CROP ROTATION

**DO NOT** plant crops in previously treated areas unless in compliance with the Rotational Restrictions found below, or in the specific crop use directions. Illegal residues may result if Rotational Restrictions are not followed. Consult **POTATO VINE** for Rotational Restrictions specifically following **Axill Solutions Glufosinate 280SL** application to potatoes as a vine desiccant.

Crop	Minimum Rotational Interval
Canola; Corn, Sweet Corn; Cotton; Soybeans; and Sugarbeets	0 days  (may be planted at any time after <b>Axill Solutions Glufosinate 280SL</b> application)
Root and Tuber Vegetables; Leafy Vegetables; Brassica Leafy Vegetables; and Small Grains (Barley, Buckwheat, Oats, Rye, Teosinte, Triticale, Wheat)	70 days
Other Crops*	180 days

\*For all crops not listed in the table above, there must be a minimum rotation interval of 180 days.

## WEED RESISTANCE MANAGEMENT

Glufosinate-ammonium, the active ingredient in this product, is a Group 10 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 10 herbicides. Weed species with acquired resistance to Group 10 herbicides may eventually dominate the weed population if Group 10 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by **Axill Solutions Glufosinate 280SL** or other Group 10 herbicides. Users must scout before and after application.

To delay herbicide resistance:

- Rotate the use of **Axill Solutions Glufosinate 280SL** or other Group 10 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout fields after application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species;
- (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seeds production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seeds.
- If a weed pest population continues to progress after treatment with this product discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management directions for specific crops and resistant weed biotypes.
- For further information or to report suspected resistance, contact a Axill Solutions, LLC (1-559-936-0444) retailer or representative.

### MANDATORY SPRAY DRIFT MANAGEMENT

- When applying to crops via aerial application equipment, the spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip or rotor blade vortices. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
- When applying to crops via aerial application equipment, applicators must use ½ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.
- For aerial applications, **DO NOT** release spray at a height greater than 10 feet above the crop canopy, unless a greater application height is necessary for pilot safety.
- For ground applications and aerial applications, select nozzle and pressure that deliver medium to coarse spray droplets as indicated in nozzle manufacturer's catalogues and in accordance with ASABE Standard 572.1.
- Spray at the appropriate boom height based on nozzle selection and nozzle spacing, but **DO NOT** exceed a boom height of 24 inches above target pest or crop canopy. Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions. Automated boom height controllers are advised with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.
- For non-crop vegetation management ground applications, apply with the nozzle height no more than 4 feet above the ground or target vegetation, unless necessitated by the application equipment. Examples would include roadside, railroad, utility rights of way, forestry, and other industrial vegetation management applications where safety or natural barriers obstruct application.

### SPRAY DRIFT ADVISORIES

#### POLLINATOR ADVISORY STATEMENT

This product contains an herbicide. Follow all label directions and precautions to minimize potential off-target exposure in order to prevent effects to non-target plants adjacent to the treated site which may serve as habitat or forage for pollinators.

#### Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See Wind, Temperature and Humidity, and Temperature Inversions sections of this label.

#### Techniques for Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures advised for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

#### Controlling Droplet Size – Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations. **AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**
- **Nozzle Type** - Solid stream nozzles (including disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - Longer booms increase drift potential. Therefore, a shorter boom length is advised.
- **Application Height** - Application more than 10 ft. above the canopy increases the potential for spray drift.

#### Boom Height

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom must remain level with the crop and have minimal bounce.

#### Drift Reduction Technology (DRT)

The EPA Drift Reduction Technology (DRT) Program was developed to encourage the manufacture, marketing, and use of spray technologies scientifically verified to significantly reduce pesticide drift. The use of DRTs should result in significantly less pesticide spray applications drifting and being deposited in areas not targeted by those applications, compared to spray technologies that **DO NOT** meet the minimum DRT standard. EPA-verified drift reduction technologies (DRTs) and their ratings will be added to the following webpage as they become available: <https://www.epa.gov/reducing-pesticide-drift/epa-verified-and-rated-drift-reduction-technologies>

#### Wind

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.** **Note:** Local terrain can influence wind patterns. Every applicator needs to be familiar with local wind patterns and how they affect spray drift.

## Temperature and Humidity

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

## Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## Shielded Sprayers

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

### USE DIRECTIONS FOR LISTED BERRY, TREE AND VINE CROPS

**Axill Solutions Glufosinate 280SL** can be applied to the following Berry, Tree and Vine orchard crops:

- Bushberry (crop subgroup 13B): blueberry, currant, elderberry, gooseberry, huckleberry
- Berries (other): lingonberry; juneberry; salal
- Citrus (crop group 10-10): lemon, orange, grapefruit, lime, mandarin, tangerine, tangelo, calamondin, kumquat, pummelo, citron, citrus hybrids, Tangor, and cultivars, varieties and/or hybrids of these
- Olives
- Pome Fruits (crop group 11-10): apples, pear, crabapple, loquat, mayhaw, quince, azarole, Medlar, Tejocote, cultivars, varieties and/or hybrids of these
- Stone Fruit (crop group 12-12): apricot, cherry, peach, nectarine, plum, capulin, jujube, Sloe and cultivars, varieties and/or hybrids of these
- Tree Nuts (crop group 14-12): almond, filberts, hickory nuts, macadamia nuts (bush nuts), pecans, pistachio, walnut
- Vine: all grape varieties (table, wine, raisin)

Application of **Axill Solutions Glufosinate 280SL** can be made via broadcast, spot or directed spray or banded spray applications

## Use Rate and Timing

**Broadcast Application:** Make broadcast application at the following use rates, depending on height of weeds or growth phase of grasses:

48 fl. oz. product/A (0.87 lb. a.i./A) – Weeds <3" in height

56 fl. oz. product/A (1.02 lbs. a.i./A) – Weeds <6" in height, pre-tiller grasses

56 - 82 fl. oz./A (1.02 - 1.5 lbs. a.i./A) – Weeds >6" in height and/or grasses that have tillered

**Spot or Directed Spray:** Make application to weeds until foliage is wet, but not to the point of runoff. Use 1.7 fl. oz. **Axill Solutions Glufosinate 280SL** (0.03 lb. a.i.) per gallon of water.

**Banded Application:** Rates indicated above are for broadcast use. The equivalents must be adjusted to reflect the actual treated area. The following formulas indicate accurate rate and volume for banded uses:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Rate per acre broadcast} = \text{Amount of herbicide needed for treatment}$$

**Sucker Control:** Apply 56 fl. oz. product/A (1.02 lbs. a.i./A) in a split application to young, green uncalled suckers that are no more than 12" in length. Wait approximately 4 weeks between applications. Sucker foliage must be thoroughly covered with product.

## Important:

- **Axill Solutions Glufosinate 280SL** can be applied as a directed broadcast spray, banded or spot treatment, to control weeds and undesirable vegetation in tree, vine and berries listed above.
- Consult **WEED LIST A** for weeds controlled.
- For best results, use appropriate rates for size of weeds, make sure spray consistently and fully covers weeds, and apply to emerged weeds that are young and actively growing and which are less than 3" in height.
- For enhanced performance of **Axill Solutions Glufosinate 280SL**, apply during warm and sunny weather, and high humidity.
- Lower use rate, unfavorable environment, or weed growth stage at time of treatment could result in regrowth of weeds. Additional application of **Axill Solutions Glufosinate 280SL** may be needed for control of weeds growing from seeds or in-ground portions of weeds.
- It is best to use highest labeled application rate for weeds in thick populations or under stress (including prior treatments with other herbicides).
- **Axill Solutions Glufosinate 280SL** solution can injure or severely damage green bark, branches or vegetation, or desirable non-target plants. Avoid contact with mixture, spray, drift, or mist. Trunks with callused, established brown bark, or shielded by nonporous wraps, grow tubes or waxed containers can be sprayed with **Axill Solutions Glufosinate 280SL**.
- Thoroughly clean application equipment following use.

## Tank Mixes

Unexposed plant parts or residual weed growth is not controlled by **Axill Solutions Glufosinate 280SL**. For residual control or control of a broader spectrum of weeds, or to support product performance, **Axill Solutions Glufosinate 280SL** can be mixed with other herbicides registered for use on berries, trees, and vine crops. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mix. When using **Axill Solutions Glufosinate 280SL** in a tank mix, additional surfactant is not required.

## Restrictions:

- **DO NOT** apply more than 164 fl. oz./A **Axill Solutions Glufosinate 280SL** (3 lbs. a.i./A) to bushberry (crop subgroup 13B) or stone fruit (crop group 12-12) within a 1-year period.
- **DO NOT** apply more than 246 fl. oz./A **Axill Solutions Glufosinate 280SL** (4.5 lbs. a.i./A) to tree nuts (crop group 14-12) and vines, pome fruit (crop group 11-10), citrus (crop group 10-10) and olives within a 1-year period.
- **DO NOT** apply more than 82 fl. oz./A (1.5 lbs. a.i./A) in a single application.
- **DO NOT** make more than 2 applications per year to bushberry (crop subgroup 13B), or stone fruit (crop group 12-12).
- **DO NOT** make more than 3 applications to tree nuts (crop group 14-12), vines, pome fruit (crop group 11-10), citrus (crop group 10-10), or olives per year.
- Pre-harvest interval is 14 days.
- **DO NOT** graze or harvest and/or feed berry, tree nut and vine orchard cover crops to livestock.
- **DO NOT** make directed spray or spot applications to vine trunk or tree trunk, or spot applications to suckers (to avoid injury).
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** apply this product using aerial application.
- **DO NOT** reapply within 14 days of previous application for berries, tree nuts, vine, citrus fruits, pome fruits and olives.
- **DO NOT** reapply within 28 days of previous application for stone fruits.
- **DO NOT** retreat after previous herbicide application until weeds have reached adequate height for effective treatment.

## WEED LIST A - For Berry, Tree and Vine Crops

Table 1

Broadleaf Weeds			
Alkali sida	Fleabane, annual	Morningglory, entireleaf	Redmaids
Ammannia, purple	Goosefoot	Morningglory, ivyleaf	Shepherd's Purse
Arrowhead, California	Gromwell, field	Morningglory, pitted	Smartweed, Pennsylvania
Buckwheat, wild	Groundcherry, cutleaf	Mullein, turkey	Sowthistle, annual
Buffalobur	Groundsel, common	Mustard, wild	Spurge, prostrate
Burclover, California	Henbit	Nettle	Starthistle, yellow
Carpetweed	Jimsonweed	Nightshade, black	Sunflower, common
Chickweed, common	Knotted	Nightshade, eastern black	Sunflower, prairie
Chinese thornapple	Kochia	Nightshade, hairy	Sunflower, volunteer
Cocklebur, common	Lambsquarters, common	Pennycress	Swinecress
Cudweed	Lettuce, miner's	Pigweed, redroot	Thistle, Russian
Cutleaf evening primrose	Lettuce, prickly	Pineapple-weed	Turnip, wild
Dodder	London rocket	Puncturevine	Velvetleaf
Eclipta	Mallow, common	Purslane, common	Vervain
Fiddleneck	Malva (little mallow)	Radish, wild	Vetch
Filaree	Marestail	Ragweed, common	Virginia copperleaf
Filaree, redstem	Mayweed	Ragweed, giant	Willowherb, panicle
Biennial and Perennial Weeds			
Aster, white heath	Dandelion	Mustard, tansy	Rose, wild
Bindweed, field	Dock, curly	Nutsedge, purple	<i>Rubus</i> spp.
Bindweed, hedge	Dogbank (hemp)	Nutsedge, yellow	Spurge, leafy
Bluegrass, Kentucky	Fescue	Onion, wild	Thistle, bull
Bromegrass, smooth	Goldenrod, gray	Orchardgrass	Thistle, Canada
Bulrush <sup>1</sup>	Guineagrass	Paragrass	Thistle, musk
Burdock	Horsetail	Plantain	Torpedograss
Clover, Alsike	Lovegrass	Poison ivy/oak	Vaseygrass
Clover, red	Mugwort	Quackgrass	Woodsorrel
Clover, white	Mullein, common	Rocket, yellow	Yarrow, common
Dallisgrass			

(continued)

**WEED LIST A - For Berry, Tree and Vine Crops**

**Table 1** (continued)

Grass Weeds			
Barnyardgrass	Crabgrass, smooth	Junglerice	Shattercane
Bluegrass, annual	Cupgrass, woolly	Oat, wild	Sprangletop
Brome, rigput	Foxtail, giant	Panicum, fall	Stinkgrass
Bromegrass, downy	Foxtail, green	Panicum, Texas	Wheat, volunteer
Canarygrass	Foxtail, yellow	Rush, toad <sup>1</sup>	Windgrass
Chess, soft	Goosegrass	Ryegrass, annual <sup>2</sup>	Witchgrass
Crabgrass, large	Johnsongrass, seedling	Sandbur, field	

<sup>1</sup>Suppression only.  
<sup>2</sup>Make application to annual ryegrass before it reaches 3" tall.

**USE DIRECTIONS FOR CANOLA**

**Axill Solutions Glufosinate 280SL** can be applied to glufosinate-resistant canola to control weeds.

**Use Rate and Timing**

Apply 22 fl. oz./A product (0.4 lb. a.i./A) to canola in cotyledon to early bolt stage. A second application of 22 fl. oz./A product (0.4 lb. a.i./A) may be used for control of later emerging weeds. Consult **WEED LIST C** (20 - 22 fl. oz./A rate) (0.37 - 0.4 lb. a.i./A rate) for weed species controlled and weed height for optimum control.

**Important:**

- For enhanced performance of **Axill Solutions Glufosinate 280SL**, apply during warm and sunny weather and high humidity
- For best results, make sure spray consistently and fully covers weeds, and apply to emerged weeds that are young and actively growing and which are less than 3" in height
- Use of **Axill Solutions Glufosinate 280SL** may result in a temporary slight discoloration of canola after application, which will not influence yield, growth, or maturity of the crop.
- Early season weed control is necessary for best harvest
- **Axill Solutions Glufosinate 280SL** will not control any volunteer glufosinate-resistant plants (corn, cotton, soybean or sugarbeets) that are left from the previous season.

If canola is injured or adversely affected by environmental stress or conditions (including excessive rainfall, fog, heavy dew, cool temperatures, cloudiness, drought, pest pressure, etc.) or prior herbicide application, weed control can be reduced

**Tank Mixes**

To support product performance, **Axill Solutions Glufosinate 280SL** can be mixed with other herbicides registered for use on canola. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mix. When using **Axill Solutions Glufosinate 280SL** in a tank mix with a grass herbicide used at a reduced rate (including a herbicide containing the active ingredients quizalofop, sethoxydim or clethodim) the AMS rate may be reduced to 1.5 lbs./A. When using **Axill Solutions Glufosinate 280SL** in a tank mix, additional surfactant is not required.

**Spray Additive**

If weeds are under stress or if more persistent weeds (including velvetleaf or lambsquarters) are present, addition of ammonium sulfate (AMS) at a rate of 1.5 - 3 lbs./A to **Axill Solutions Glufosinate 280SL** can enhance results. Choose appropriate rate based on temperatures and environmental conditions, the probability of leaf burn, or tank mixes. If necessary, other additives, including drift control agents can be used. Use of an antifoam agent is suggested. Use care when adding additional crop oils or surfactants, as the risk of an adverse crop response can be increased.

**Restrictions:**

- **DO NOT** apply more than 22 fl. oz./A (0.4 lb. a.i./A) per application.
- **DO NOT** make more than 2 applications of **Axill Solutions Glufosinate 280SL** per year.
- **DO NOT** exceed the max yearly application rate of 44 fl. oz./A (0.8 lb. a.i./A).
- **DO NOT** reapply within 10 days of previous applications.
- **DO NOT** make additional (post-emergent) applications to canola when **Axill Solutions Glufosinate 280SL** is used as a burndown prior to planting.
- **DO NOT** apply within 65 days of harvest.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** hay or graze treated canola.
- **DO NOT** apply **Axill Solutions Glufosinate 280SL** to glufosinate-resistant canola in the following states: AL, DE, GA, KY, MD, NJ, NC, SC, TN, VA, and WV.

**USE DIRECTIONS FOR BURNDOWN - CANOLA**

**Axill Solutions Glufosinate 280SL** can be applied prior to planting or emergence of conventional or transgenic canola to act as a burndown agent for existing weed species.

**Axill Solutions Glufosinate 280SL** can be applied via 2 different use patterns. Timing 1 use pattern is used for low to medium weed pressure, and when weeds are small and actively growing. Timing 2 use pattern is used if weed pressure is high and weeds are larger and actively growing.

**Table 2: RESTRICTIONS - Burndown Use Rates in Canola**

Use Pattern	Use Rate: Burndown	Use Rate: In Season	Cumulative Maximum use rate per year
TIMING 1	29 fl. oz./A (0.53 lb. a.i./A)	None	36 fl. oz./A (0.66 lb. a.i./A)
TIMING 2	Up to 36 fl. oz./A (0.66 lb. a.i./A)	None	36 fl. oz./A (0.66 lb. a.i./A)

Apply at least 29 fl. oz./A of **Axill Solutions Glufosinate 280SL** (0.53 lb. a.i./A) just before planting or emergence. If weeds are large or weed pressure is high, due to environmental conditions preventing timely use of **Axill Solutions Glufosinate 280SL**, a single application of up to 36 fl. oz./A (0.66 lb. a.i./A) can be made.

**Restrictions:**

- **DO NOT** apply more than 36 fl. oz./A (0.66 lb. a.i./A) per burndown application.
- **DO NOT** apply more than 36 fl. oz./A (0.66 lb. a.i./A) per year.
- **DO NOT** make more than 1 burndown application per year.
- **DO NOT** make additional (post-emergent) applications to canola when **Axill Solutions Glufosinate 280SL** is used as a burndown prior to planting.

**USE DIRECTIONS FOR SWEET CORN\*, FIELD CORN AND SILAGE CORN**

**\*Not for use on sweet corn in California.**

**Axill Solutions Glufosinate 280SL** can be applied to glufosinate-resistant corn to control weeds.

**Use Rate and Timing**

Apply 22 fl. oz./A product (0.4 lb. a.i./A) to field and silage corn, or 20 fl. oz./A product (0.37 lb. a.i./A) to sweet corn. A second application of 22 fl. oz. (field and silage corn) (0.4 lb. a.i./A) or 20 fl. oz. (sweet corn) (0.37 lb. a.i./A), or tank mix with a residual herbicide may be used for control of later emerging weeds. Consult **WEED LIST C** (20 - 22 fl. oz. rate) (0.37 - 0.4 lb. a.i./A rate) for weed species controlled and optimum weed height for control.

**Timing**

- Corn up to 24" tall or V7 stage of growth (7 developed collars), whichever comes first – Apply over the top or ground application with drop nozzle.
- Corn 24" to 36" tall – Ground application with drop nozzle (avoid spraying corn stalk leaf axis or whorls).

**Important:**

- For best results, make sure spray consistently and fully covers weeds and apply to emerged weeds that are young and actively growing and which are less than 3" in height.
- For enhanced performance of **Axill Solutions Glufosinate 280SL**, apply during warm and sunny weather and high humidity.
- Early season weed control is necessary for best harvest.

If corn is injured or adversely affected by environmental stress or conditions (including excessive rainfall, fog, heavy dew, cool temperatures, cloudiness, drought, pest pressure, etc.) or prior herbicide applications, weed control can be reduced.

**Tank Mixes:** To support product performance, **Axill Solutions Glufosinate 280SL** can be mixed with other herbicides registered for use on corn (sweet, field or silage). It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mix. Additional surfactant is not needed when tank mixing. If tank mixing with products containing the active ingredients carfentrazone-ethyl, metolachlor, s-metolachlor, and products with a combination of s-metolachlor, mesotrione and atrazine, use half the rate with **Axill Solutions Glufosinate 280SL**, to reduce the risk for adverse crop response. If tank mixing with products containing pendimethalin, be aware that reduced control of barnyardgrass, fall panicum, field sandbur, yellow foxtail and volunteer corn can occur.

**Spray Additive:** If weeds are under stress or if more persistent weeds (including velvetleaf or lambsquarters) are present, addition of ammonium sulfate (AMS) at a rate of 1.5 - 3 lbs./A to **Axill Solutions Glufosinate 280SL** can enhance results. Choose appropriate rate based on temperatures and environmental conditions, the probability of leaf burn, or tank mixes. Use of an antifoam agent is suggested. Use care when adding additional crop oils or surfactants, as the risk of an adverse crop response can be increased.

**Restrictions:**

- **DO NOT** apply more than 22 fl. oz./A (0.4 lb. a.i./A) per application for field and silage corn.
- **DO NOT** apply more than 20 fl. oz./A (0.37 lb. a.i./A) per application for sweet corn.
- **DO NOT** more than 2 applications of **Axill Solutions Glufosinate 280SL** to corn per year.
- **DO NOT** exceed the maximum yearly application rate of 44 fl. oz. (0.8 lb. a.i./A) for field, silage.
- **DO NOT** exceed the maximum yearly application rate of 40 fl. oz. (0.8 lb. a.i./A) for sweet corn.
- **DO NOT** apply within 10 days of previous application.

- When **Axill Solutions Glufosinate 280SL** is used as a burndown prior to planting corn, no additional (post-emergent) applications can be made to corn.
- Pre-harvest interval for sweet corn ears: 50 days
- Pre-harvest interval for sweet corn stover: 55 days
- Pre-harvest interval for field and silage corn forage: 60 days
- Pre-harvest interval for field and silage corn grain and fodder: 70 days
- **DO NOT** apply through any type of irrigation system.
- Nitrogen solutions are not to be used as spray carriers.

#### USE DIRECTIONS FOR BURNDOWN – CORN (FIELD, SILAGE AND SWEET)

**Axill Solutions Glufosinate 280SL** can be applied prior to planting or emergence of conventional or transgenic corn to act as a burndown agent for existing weed species.

**Axill Solutions Glufosinate 280SL** can be applied via 2 different use patterns. Timing 1 use pattern is used for low to medium weed pressure, and when weeds are small and actively growing. Timing 2 use pattern is used if weed pressure is high and weeds are larger and actively growing.

**Table 3: RESTRICTIONS - Burndown Use Rates in Corn**

Use Pattern	Use Rate: Burndown	Use Rate: In Season	Cumulative Maximum use rate per year
TIMING 1	29 fl. oz./A (0.53 lb. a.i./A)	None	36 fl. oz./A (0.66 lb. a.i./A)
TIMING 2	Up to 36 fl. oz./A (0.66 lb. a.i./A)	None	36 fl. oz./A (0.66 lb. a.i./A)

Apply at least 29 fl. oz./A of **Axill Solutions Glufosinate 280SL** (0.53 lb. a.i./A) just before planting or emergence. If weeds are large or weed pressure is high, due to environmental conditions preventing timely use of **Axill Solutions Glufosinate 280SL**, a single application of up to 36 fl. oz./A (0.66 lb. a.i./A) can be made.

#### Restrictions:

- **DO NOT** apply more than 36 fl. oz./A (0.66 lb. a.i./A) per burndown application.
- **DO NOT** apply more than 36 fl. oz./A (0.66 lb. a.i./A) per year.
- **DO NOT** make more than 1 burndown application per year.
- When **Axill Solutions Glufosinate 280SL** is used as a burndown prior to planting corn, no additional (post-emergent) applications can be made to corn.

#### USE DIRECTIONS FOR COTTON

**Axill Solutions Glufosinate 280SL** can be applied to glufosinate-resistant cotton (broadcast, over-the-top post-emergence or directed spray) to control weeds, or conventional (non-glufosinate-resistant) cotton (post-emergence hooded/shielded spray) to control weeds. If **Axill Solutions Glufosinate 280SL** comes in contact with non-glufosinate-resistant cotton plants (foliage or stems), serious injury or loss of plant could occur. **Axill Solutions Glufosinate 280SL** can also be used for post-harvest applications.

#### Use Rate and Timing

**Axill Solutions Glufosinate 280SL** can be applied to cotton via 2 different use patterns. Timing 1 use pattern is used if weeds are particularly large, or weed pressure is high due to environmental conditions preventing timely use of **Axill Solutions Glufosinate 280SL**. Timing 2 use pattern is used for low to medium weed pressure. Consult **WEED LIST C** for weed species controlled.

**Banded Application:** Rates indicated above are for broadcast use. The equivalents must be adjusted to reflect the actual treated area. The following formulas indicate accurate rate and volume for banded uses:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per acre} = \text{Amount of banded product needed per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast spray volume per acre} = \text{Banded spray volume needed per acre}$$

**Glufosinate-Resistant Cotton:** Apply product foliarly over the top or directed spray to the lower 1/3 of the cotton stand.

#### Non-Glufosinate-Resistant Cotton - Hooded Spray Application:

Apply **Axill Solutions Glufosinate 280SL** via a hooded sprayer that is designed to direct product spray to the weeds and minimize exposure to cotton plants. When using this method of application, take all possible steps to avoid exposing cotton plants to product spray. Key points for using hooded sprayer are:

- A hooded sprayer operates so that the top and sides are enclosed by a hood, protecting cotton from product spray.
- Set up hooded sprayer so that it is run on or skims across the field, and operate the sprayer in a manner and speed that keeps the hood from bouncing or raising off the ground. (**Note:** If hood becomes raised, it can allow spray particles to escape, which could cause damage or loss of cotton plant.)
- Adjust hoods to protect desirable plants and use nozzles that deliver uniform exposure.
- Avoid applying where ground is sloped or uneven, and could allow spray hoods to be raised off the ground.

### Important:

- For best results, make sure spray consistently and fully covers weeds, and apply to emerged weeds that are young and actively growing and which are less than 3" in height.
- For enhanced performance of **Axill Solutions Glufosinate 280SL**, apply during warm and sunny weather, and high humidity.
- If cotton are injured or adversely affected by environmental stress or conditions (including excessive rainfall, fog, heavy dew, cool temperatures, cloudiness, drought, pest pressure, etc.) or prior herbicide application, weed control can be reduced.
- Consult **WEED CHART B** for weed control, and if a mixed population of weeds are present, use the rate necessary to control all weed species.
- Additional application of **Axill Solutions Glufosinate 280SL** or tank mix with other herbicide may be needed for control of weeds growing from seeds or in-ground portions of weeds.
- Early season weed control is necessary for best harvest.
- **Axill Solutions Glufosinate 280SL** will not control any volunteer glufosinate-resistant plants (corn, cotton, soybean, sugarbeet) that are left from the previous season.

### Tank Mixes

To support product performance, **Axill Solutions Glufosinate 280SL** can be mixed with other herbicides registered for use on cotton. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mix.

### Spray Additive

If weeds are under stress or if more persistent weeds (including velvetleaf or lambsquarters) are present, addition of ammonium sulfate (AMS) at a rate of 1.5 - 3 lbs./A to **Axill Solutions Glufosinate 280SL** can enhance results. Choose appropriate rate based on temperatures and environmental conditions, the probability of leaf burn, or tank mixes. Use of an antifoam agent is suggested. Use care when adding additional crop oils or surfactants, as the risk of an adverse crop response can be increased.

### Restrictions:

- **DO NOT** apply more than 43 fl. oz./A (0.79 lb. a.i./A) in a single application for post-harvest burndown.
- **DO NOT** make more than 3 applications per year of **Axill Solutions Glufosinate 280SL** at a rate of 29 fl. oz./A (0.53 lb. a.i./A).
- If the single application rate exceeds 29 fl. oz./A (0.53 lb. a.i./A) then the yearly maximum use rate is 72 fl. oz./A (1.32 lbs. a.i./A) (including all application timings).
- If applying a maximum application rate of 29 fl. oz./A (0.53 lb. a.i./A), the maximum yearly application rate is not to exceed 87 fl. oz./A (1.6 lbs. a.i./A), including all application timings.
- If large weeds or dense infestation is present, because timely application was not possible (due to environmental conditions), an application rate not to exceed 43 fl. oz./A (0.79 lb. a.i./A) can be made, and a second application may be made at a maximum rate of 29 fl. oz./A (0.53 lb. a.i./A).
- If applying a maximum application rate greater than 29 fl. oz./A (0.53 lb. a.i./A), the maximum yearly application rate is not to exceed 72 fl. oz./A (1.32 lbs. a.i./A), including all application timings.
- **Post-Harvest Application:** When using **Axill Solutions Glufosinate 280SL** as a post-harvest burndown treatment to cotton fields, a single application not to exceed 43 fl. oz./A (0.79 lb. a.i./A) can be applied. Adjust use rate to correspond with size of weeds.
- **DO NOT** reapply within 10 days of prior application.
- Pre-harvest interval is 70 days.
- **DO NOT** apply through any type of irrigation system.
- **Axill Solutions Glufosinate 280SL** cannot be applied to cotton south of Tampa in Florida (Route 60) or in Hawaii (except for application to breeding nurseries or test plots).

**Table 4: RESTRICTIONS – Broadcast, Over-the-Top Post-Emergence, or Directed Spray Use Rates in Cotton**

Use Pattern	Use Rate Application 1	Use Rate Application 2	Use Rate Application 3	Cumulative Maximum use rate per year
TIMING 1	32 - 43 fl. oz./A (0.59 - 0.79 lb. a.i./A)	29 fl. oz./A (0.53 lb. a.i./A)	None	72 fl. oz./A (1.32 lbs. a.i./A)
TIMING 2	29 fl. oz./A (0.53 lb. a.i./A)	29 fl. oz./A (0.53 lb. a.i./A)	29 fl. oz./A (0.53 lb. a.i./A)	87 fl. oz./A (1.59 lbs. a.i./A)

### USE DIRECTIONS FOR BURNDOWN - COTTON

**Axill Solutions Glufosinate 280SL** can be applied prior to planting or emergence of conventional or transgenic cotton to act as a burndown agent for existing weed species.

**Axill Solutions Glufosinate 280SL** can be applied via 2 different use patterns. Timing 1 use pattern is used for low to medium weed pressure, and when weeds are small and actively growing. Timing 2 use pattern is used if weed pressure is high and weeds are larger and actively growing.

**Table 5: RESTRICTIONS - Burndown Use Rates in Cotton**

Use Pattern	Use Rate: Burndown	Use Rate: In Season	Cumulative Maximum Use Rate per Year
TIMING 1	29 fl. oz./A (0.53 lb. a.i./A)	22 - 29 fl. oz./A (2 applications)* (0.4 - 0.53 lb. a.i./A)	87 fl. oz./A (1.59 lbs. a.i./A)
TIMING 2	30 - 43 fl. oz./A (0.55 - 0.79 lb. a.i./A)	22 - 29 fl. oz./A (1 application)* (0.4 - 0.53 lb. a.i./A)	72 fl. oz./A (1.32 lbs. a.i./A)

\*In season application is made via hooded sprayer in non-glufosinate-resistant cotton. Apply at least 29 fl. oz./ of **Axill Solutions Glufosinate 280SL** (0.53 lb. a.i./A) just before planting or emergence. If weeds are large or weed pressure is high, due to environmental conditions preventing timely use of **Axill Solutions Glufosinate 280SL**, a single application of up to 43 fl. oz./A (0.79 lb. a.i./A) can be made. Yearly maximum use rate is 72 fl. oz./A (1.32 lbs. a.i./A) (including all application timings) when a single application higher than 29 fl. oz./A (0.53 lb. a.i./A) is made.

**Restrictions:**

- **DO NOT** apply more than 43 fl. oz./A (0.79 lb. a.i./A) per burndown application.
- **DO NOT** apply more than 43 fl. oz./A (0.79 lb. a.i./A) per year when applying a preplant burndown application.
- **DO NOT** make more than 1 burndown application per year.

**Important:**

- For best results, make sure spray consistently and fully covers weeds, and apply to emerged weeds that are young and actively growing.
- For enhanced performance of **Axill Solutions Glufosinate 280SL**, apply during warm and sunny weather, and high humidity.

**USE DIRECTIONS FOR SOYBEANS**

**Axill Solutions Glufosinate 280SL** can be applied to glufosinate-resistant soybeans to control weeds.

**Use Rate and Timing**

Apply 22 - 29 fl. oz./A product (0.4 - 0.53 lb. a.i./A) to soybeans, from emergence up to just before bloom. If weed pressure is high, due to environmental conditions, preventing a timely use of **Axill Solutions Glufosinate 280SL**, a single application of up to 36 fl. oz./A (0.66 lb. a.i./A) can be made to soybeans beyond the V3 - V4 vegetative growth phase. A second application of 29 fl. oz./A (0.53 lb. a.i./A) can be used for later emerging weeds. Consult **WEED LIST C** for appropriate application rate based on weed type and size.

**Important:**

- For enhanced performance of **Axill Solutions Glufosinate 280SL**, apply during warm and sunny weather and high humidity.
- For best results, make sure spray consistently and fully covers weeds, and apply to emerged weeds that are young and actively growing and which are less than 3" in height.
- Early season weed control is necessary for best harvest.
- **Axill Solutions Glufosinate 280SL** can provide complete weed control when applied in a timely manner, however tank mixing or use of **Axill Solutions Glufosinate 280SL** in conjunction with residual herbicides can help with reduction of early season weed competition, and control of later emerging weeds (particularly if environmental conditions prevented timely application of **Axill Solutions Glufosinate 280SL**).
- **Axill Solutions Glufosinate 280SL** will not control any volunteer glufosinate-resistant plants (corn, cotton, soybean, sugarbeet) that are left from the previous season.

If soybeans are injured or adversely affected by environmental stress or conditions (including excessive rainfall, fog, heavy dew, cool temperatures, cloudiness, drought, pest pressure, etc.) or prior herbicide application, weed control can be reduced.

**Tank Mixes**

To support product performance, **Axill Solutions Glufosinate 280SL** can be mixed with other herbicides registered for use on soybeans. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mix. Additional surfactant is not necessary when tank mixing.

**Spray Additives**

If weeds are under stress or if more persistent weeds (including velvetleaf or lambsquarters) are present, addition of ammonium sulfate (AMS) at a rate of 1.5 - 3 lbs./A to **Axill Solutions Glufosinate 280SL** can enhance results. Use of an antifoam agent is suggested. Use care when adding additional crop oils or surfactants, as the risk of an adverse crop response can be increased.

**Restrictions:**

- **DO NOT** make more than 2 applications of **Axill Solutions Glufosinate 280SL** at above rates to soybeans per year (including preplant burndown).
- **DO NOT** exceed the max yearly application rate of 65 fl. oz./A (1.2 lbs. a.i./A).
- **DO NOT** apply more than 36 fl. oz./A (0.66 lb. a.i./A) in a single application.
- **DO NOT** apply within 10 days of previous application.
- Pre-harvest interval is 70 days.
- **Axill Solutions Glufosinate 280SL** is not to be applied through any type of irrigation system.
- Soybeans treated with **Axill Solutions Glufosinate 280SL** cannot be cut for hay or grazed.
- Nitrogen solutions are not to be used as spray carriers.

## USE DIRECTIONS FOR BURNDOWN - SOYBEANS

**Axill Solutions Glufosinate 280SL** can be applied prior to planting or emergence of conventional or transgenic soybeans to act as a burndown agent for existing weed species.

**Axill Solutions Glufosinate 280SL** can be applied via 2 different use patterns. Timing 1 use pattern is used for low to medium weed pressure, and when weeds are small and actively growing. Timing 2 use pattern is used if weed pressure is high and weeds are larger and actively growing.

**Table 6: RESTRICTIONS – Burndown Use Rates in Soybeans**

Use Pattern	Use Rate: Burndown	Use Rate: In Season	Cumulative Maximum use rate per year
TIMING 1	29 fl. oz./A (0.53 lb. a.i./A)	22 - 29 fl. oz./A (0.4 - 0.53 lb. a.i./A)	65 fl. oz./A (1.2 lbs. a.i./A)
TIMING 2	Up to 36 fl. oz./A (0.66 lb. a.i./A)	22 - 29 fl. oz./A (0.4 - 0.53 lb. a.i./A)	65 fl. oz./A (1.2 lbs. a.i./A)

Apply at least 29 fl. oz./A of **Axill Solutions Glufosinate 280SL** (0.53 lb. a.i./A) just before planting or emergence. If weeds are large or weed pressure is high, due to environmental conditions preventing timely use of **Axill Solutions Glufosinate 280SL**, a single application of up to 36 fl. oz./A (0.66 lb. a.i./A) can be made. For soybeans, an additional in-season application up to 29 fl. oz./A (0.53 lb. a.i./A) can be made.

### Restrictions:

- **DO NOT** apply more than 36 fl. oz./A (0.66 lb. a.i./A) per burndown application.
- **DO NOT** apply more than 36 fl. oz./A (0.66 lb. a.i./A) per year preplant burndown.
- **DO NOT** make more than 1 burndown application per year.

## USE DIRECTIONS FOR SUGARBEETS

(Not for use in California)

**Axill Solutions Glufosinate 280SL** can be applied to glufosinate-resistant sugarbeets to control weeds.

### Use Rate and Timing

Apply up to 30 fl. oz./A product (0.55 lb. a.i./A) to sugarbeets, from cotyledon stage up to 10-leaf stage. A second application of up to 30 fl. oz./A (0.55 lb. a.i./A) will be needed to control later emerging weeds. Consult **WEED LIST B** for weed species controlled. Optimum control is obtained when weeds are 1 inch or less in height or diameter.

### Important:

- To avoid reduced performance of **Axill Solutions Glufosinate 280SL**, **DO NOT** apply when heavy dew, fog, or mist/rain are present.
- For best results, make sure spray consistently and fully covers weeds, and apply to emerged weeds that are young and actively growing and which are less than 3" in height.
- For best results, make a cultivation 5 days before or 5 days after application of **Axill Solutions Glufosinate 280SL**.
- Early season weed control is necessary for best harvest.
- If sugarbeets are injured or adversely affected by environmental stress or conditions (including excessive rainfall, fog, heavy dew, cool temperatures, cloudiness, drought, pest pressure, etc.) or prior herbicide application, weed control can be reduced.
- Weeds that emerge after **Axill Solutions Glufosinate 280SL** has been applied will not be controlled, as **Axill Solutions Glufosinate 280SL** does not exhibit residual activity. If crop experiences a rain event within 4 hours of application, retreatment may be necessary, however after 4 hours, **Axill Solutions Glufosinate 280SL** is rainfast.

### Tank Mixes

To support product performance, **Axill Solutions Glufosinate 280SL** can be mixed with other herbicides registered for use on sugarbeets. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mix.

### Spray Additives

If necessary, drift control agents can be used with **Axill Solutions Glufosinate 280SL**. **DO NOT** use surfactants. If weeds are under stress or if more persistent weeds (including velvetleaf or lambsquarters) are present, addition of ammonium sulfate (AMS) at a rate of 1.5 - 3 lbs./A to **Axill Solutions Glufosinate 280SL** can enhance results. Choose appropriate rate based on temperatures and environmental conditions, the probability of leaf burn, or tank mixes. Use of an antifoam agent is suggested.

### Restrictions:

- **DO NOT** make more than 2 applications per year of **Axill Solutions Glufosinate 280SL**.
- **DO NOT** apply with 10 days of previous application.
- **DO NOT** exceed the max yearly application rate of 60 fl. oz./A (1.1 lbs. a.i./A).
- **DO NOT** apply more than 30 fl. oz./A (0.55 lb. a.i./A) in a single application.
- Pre-harvest interval is 60 days.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** hay or graze sugarbeets treated with product.
- Rotation crop plantback in sugarbeet field treated with **Axill Solutions Glufosinate 280SL** is 120 days after last application for all crops except wheat, barley, buckwheat, millet, oats, rye, sorghum and triticale (70-day plantback restriction) or corn, cotton, soybeans, canola and sugar beets glufosinate-resistant crops (0-day plantback restriction).

**WEED LIST B - For Sugar Beets**

**Table 7** - The weed table indicates rates of product to be used for control of weeds based on weed height. If weed population consists of mixed species, apply rate indicated that will be efficacious for all species.

<b>Grass Weeds</b>			
<b>Weed Species</b>	<b>Maximum Weed Height*</b>		<b>Use Notes</b>
	<b>15 fl. oz. A (0.9 pt./A) (0.27 lb. a.i./A)</b>	<b>20 fl. oz. A (1.25 pts./A) (0.37 lb. a.i./A)</b>	
Barley, volunteer	1-2 leaf (2")	3 leaf (3")	Multiple applications may be required
Barnyard grass	1-3 leaf (2")	4-5 leaf (3")	Maximum of 1 tiller
Corn, volunteer	1-2 leaf (3")	3-4 leaf (6")	
Crabgrass, large	1-3 leaf (2")	4-5 leaf (3")	Maximum of 1 tiller
Crabgrass, smooth	1-3 leaf (2")	4-5 leaf (3")	Maximum of 1 tiller
Crabgrass, wooly	1-5 leaf (4")	(8")	
Foxtail, giant	1-4 leaf (3")	5-6 leaf (4")	Maximum of 2 tillers
Foxtail, green	1-4 leaf (3")	5-6 leaf (4")	Maximum of 2 tillers
Foxtail, yellow	1-3 leaf (1")	4 leaf (2")	Apply prior to tillering
Millet, volunteer proso	1-3 leaf (2")	4-5 leaf (3")	Max of 1 tiller
Millet, wild proso	1-3 leaf (2")	4-5 leaf (3")	Max of 1 tiller
Oat, wild	1-2 leaf (2")	3 leaf (3")	Max of 1 tiller
Panicum, fall	1-3 leaf (2")	4-5 leaf (3")	Max of 1 tiller
Panicum, Texas	1-3 leaf (2")	4-5 leaf (3")	Max of 1 tiller
Sandbur, field	--	1-4 leaf (2")	Apply prior to tillering
Wheat, volunteer	1-2 leaf (2")	3 leaf (3")	Maximum of 1 tiller
<b>Perennial Weeds</b>			
Quackgrass	--	1-3 leaf (3")	Multiple applications required
Sowthistle, perennial	--	1-4 leaf (3")	Multiple applications required
Thistle, Canada	--	1-4 leaf (3")	Multiple applications required
<b>Broadleaf Weeds</b>			
Buckwheat, wild	1-4 leaf (2")	5-6 leaf (3")	
Buffalobur	1-4 leaf (2")	5-6 leaf (3")	
Carpetweed	--	1-4 leaf (2")	
Chickweed, common	1-4 leaf (2")	5-6 leaf (3")	
Cocklebur, common	1-6 leaf (3")	7-8 leaf (5")	
Kochia	(1")	(2")	
Ladysthumb	1-2 leaf (1")	3-4 leaf (3")	
Lambsquarters, common	1-2 leaf (1")	4-5 leaf (3")	
Mallow, Venice	1-4 leaf (2")	5-6 leaf (3")	
Marshelder	1-2 leaf (1")	3-4 leaf (2")	
Mustard, wild	1-4 leaf (2")	5-6 leaf (3")	
Nightshade, eastern black	1-4 leaf (2")	5-6 leaf (3")	
Pigweed, prostrate	(1")	(3")	
Pigweed, redroot	1-2 leaf (1")	3-4 leaf (3")	
Pigweed, spiny	1-2 leaf (1")	3-4 leaf (3")	
Purslane, common	(1")	(2")	
Ragweed, common	1-6 leaf (3")	7-8 leaf (5")	
Ragweed, giant	1-4 leaf (2")	5-6 leaf (3")	
Shepherd's purse	1-4 leaf (2")	5-6 leaf (3")	
Smartweed, Pennsylvania	1-2 leaf (1")	3-4 leaf (3")	
Sowthistle, annual	1-4 leaf (2")	5-6 leaf (3")	

*(continued)*

<b>Broadleaf Weeds (continued)</b>			
<b>Weed Species</b>	<b>Maximum Weed Height*</b>		<b>Use Notes</b>
	<b>15 fl. oz. A (0.9 pt./A) (0.27 lb. a.i./A)</b>	<b>20 fl. oz. A (1.25 pts./A) (0.37 lb. a.i./A)</b>	
Sunflower, common	1-6 leaf (3")	7-8 leaf (5")	
Thistle, Russian	(1")	(2")	
Velvetleaf	1-2 leaf (1")	3-4 leaf (3")	

\*Up to 30 fl. oz./A (0.55 lb. a.i./A) can be applied if weeds are taller than indicated in table.

Tank mix with herbicides containing clothodim, quizalofop, rimsulfuron or sethoxydim to enhance control of heavy populations or taller growth stages of volunteer barley, yellow foxtail, wild oats, or volunteer wheat.

#### USE DIRECTIONS FOR BURNDOWN - SUGARBEETS

**Axill Solutions Glufosinate 280SL** can be applied prior to planting or emergence of conventional or transgenic sugarbeet, to act as a burndown agent for existing weed species.

**Axill Solutions Glufosinate 280SL** can be applied via 2 different use patterns. Timing 1 use pattern is used for low to medium weed pressure, and when weeds are small and actively growing. Timing 2 use pattern is used if weed pressure is high and weeds are larger and actively growing.

**Table 8: RESTRICTIONS – Burndown Use Rates in Sugarbeets**

<b>Use Pattern</b>	<b>Use Rate: Burndown</b>	<b>Use Rate: In Season</b>	<b>Cumulative Maximum use rate per year</b>
TIMING 1	29 fl. oz./A (0.53 lb. a.i./A)	None	36 fl. oz./A (0.66 lb. a.i./A)
TIMING 2	Up to 36 fl. oz./A (0.66 lb. a.i./A)	None	36 fl. oz./A (0.66 lb. a.i./A)

Apply at least 29 fl. oz./A of **Axill Solutions Glufosinate 280SL** (0.53 lb. a.i./A) just before planting or emergence. If weeds are large or weed pressure is high, due to environmental conditions preventing timely use of **Axill Solutions Glufosinate 280SL**, a single application of up to 36 fl. oz./A (0.66 lb. a.i./A) can be made.

#### Restrictions:

- **DO NOT** apply more than 36 fl. oz./A (0.66 lb. a.i./A) per burndown application.
- **DO NOT** apply more than 36 fl. oz./A (0.66 lb. a.i./A) per year for preplant burndown.
- **DO NOT** make more than 1 burnout application per year

#### USE DIRECTIONS FOR FALLOW FIELDS

**Axill Solutions Glufosinate 280SL** can be applied to fallow fields to control or suppress weeds, as a substitute for tillage in fallow fields. See **WEED LIST C** for a list of targeted weed species.

#### Use Rate and Timing

Consult **WEED LIST C** for appropriate application rate based on weed type and size. Use 22 - 29 fl. oz./A (0.4 - 0.53 lb. a.i./A) on fallow fields for control of target weed species. Mix **Axill Solutions Glufosinate 280SL** with ammonium sulfate for application to fallow fields.

#### Tank Mixes

To support product performance, **Axill Solutions Glufosinate 280SL** can be mixed with other herbicides registered for use in fallow fields. 2,4-D, atrazine, or glyphosate can be used as tank mix partners. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mix.

When using **Axill Solutions Glufosinate 280SL** in a tank mix, additional surfactant is not required.

#### Restrictions:

- **DO NOT** apply more than 29 fl. oz./A (0.53 lb. a.i./A) of **Axill Solutions Glufosinate 280SL** per year.
- **DO NOT** apply more than 29 fl. oz./A (0.53 lb. a.i./A) in a single application.
- **DO NOT** apply more than 1 application per year.

**WEED LIST C - For Canola, Cotton, Corn (Sweet, Field, and Popcorn), Soybeans, and Fallow Fields**

**Table 9** - The weed table indicates rates of product to be used for control of weeds based on weed height.

<b>Broadleaf Weeds</b>					
Weed Species	Maximum Weed Height or Diameter (Inches)		Weed Species	Maximum Weed Height or Diameter (Inches)	
	20 - 22 fl. oz./A (0.37 - 0.4 lb. a.i./A)	29 fl. oz./A (0.53 lb. a.i./A)		20 - 22 fl. oz./A (0.37 - 0.4 lb. a.i./A)	29 fl. oz./A (0.53 lb. a.i./A)
Amaranth, Palmer	3	4	Morningglory, sharpod	2	4
Anoda, spurred	3	5	Morningglory, smallflower	4	6
Beggarweed, Florida	4	5	Morningglory, tall	6	8
Black medic	5	7	Mustard, wild	4	6
Blueweed, Texas	5	7	Nightshade, black	4	6
Buckwheat, wild	6	7	Nightshade, eastern black	6	8
Buffalobur	6	7	Nightshade, hairy	6	8
Burcucumber	6	10	Pennycress (stinkweed)	4	6
Catchweed bedstraw (cleavers)	2	4	Pigweed, redroot	3	4
Carpetweed	4	6	Pigweed, prostrate	3	4
Chickweed, common	6	8	Pigweed, spiny	3	4
Cocklebur, common	6	14	Pigweed, smooth	3	4
Copperleaf, hophornbeam	4	6	Pigweed, tumble	3	4
Cotton, volunteer	6	8	Puncturevine	4	6
Croton, tropic	3	5	Purslane, common	2	4
Croton, woolly	2	4	Pusley, Florida	*	3
Eclipta	4	6	Ragweed, common	6	10
Devil's claw	2	4	Ragweed, giant	6	12
Fleabane, annual	6	8	Senna coffee	4	6
Galinsoga, hairy	6	8	Sesbania, hemp	6	8
Galinsoga, small flower	6	7	Shepherd's Purse	6	8
Groundcherry, cutleaf	4	5	Sicklepod (java bean)	4	6
Geranium, cutleaf	4	6	Sida, prickly	4	5
Hempnettle	4	6	Smartweed, Pennsylvania	6	14
Horsenettle, Carolina <sup>1</sup>	2	4	Smellmelon	4	6
Jimsonweed	6	10	Sowthistle, annual	6	8
Knotweed	3	5	Soybeans, volunteer	6	8
Kochia	4	6	Spurge, prostrate	2	4
Ladysthumb	6	14	Spurge, spotted	2	4
Lambsquarters, common	4	6	Starbur, bristly	4	6
Mallow, common	4	6	Sunflower, common	6	14
Mallow, Venice	6	8	Sunflower, prairie	3	5
Marestail <sup>1</sup>	*	6-12	Sunflower, volunteer	6	10
Marshelder, annual	4	6	Thistle, Russian <sup>1</sup>	S	6-12
Morningglory, entireleaf	6	8	Velvetleaf	3	4
Morningglory, ivyleaf	6	8	Waterhemp, common	4	5
Morningglory, pitted	6	8	Waterhemp, tall	4	5

\*Suppression only.

<sup>1</sup>For control, a sequential application may be needed.

### Biennial and Perennial Weeds

For control of the biennial and perennial weeds listed below, apply tank mix partners or sequential applications of **Axill Solutions Glufosinate 280SL** (22 fl. oz./A followed by 22 fl. oz./A). (22 fl. oz./A = 0.4 lb. a.i./A)

Alfalfa	Bursage, woolyleaf	Johnsongrass, rhizome	Poinsettia, wild
Artichoke, Jerusalem	Chickweed, mouse-ear	Milkweed, common*	Pokeweed
Bermudagrass	Clover, Alsike	Milkweed, honeyvine*	Quackgrass*
Bindweed, field	Clover, red	Muhly, wirestem*	Sowthistle, perennial
Bindweed, hedge	Dandelion	Nightshade, silverleaf	Thistle, bull
Bluegrass, Kentucky	Dock, smooth	Nutsedge, purple*	Thistle, Canada
Blueweed, Texas	Dogbane, hemp*	Nutsedge, yellow*	Timothy*
Bromegrass, smooth	Goldenrod, gray*	Orchardgrass	Wormwood, biennial
Burdock			

\*Suppression only.

### Grass Weeds

Weed Species	Maximum Weed Height or Diameter (Inches)		Weed Species	Maximum Weed Height or Diameter (Inches)	
	20 - 22 fl. oz./A (0.37 - 0.4 lb. a.i./A)	29 fl. oz./A (0.53 lb. a.i./A)		20 - 22 fl. oz./A (0.37 - 0.4 lb. a.i./A)	29 fl. oz./A (0.53 lb. a.i./A)
Barley, volunteer <sup>3</sup>	3	4	Millet, wild-proso	6	7
Barnyardgrass	3	5	Millet, proso volunteer	6	7
Bluegrass, annual	3	5	Oat, wild <sup>2</sup>	3	4
Corn, volunteer <sup>1</sup>	10	12	Panicum, fall	3	5
Crabgrass, large <sup>2</sup>	3	5	Panicum, Texas	4	6
Crabgrass, smooth <sup>2</sup>	3	5	Rice, red	4	6
Cupgrass, woolly	6	12	Rice, volunteer <sup>1</sup>	4	6
Foxtail, bristly	6	8	Sandbur, field <sup>3</sup>	*	2
Foxtail, giant	6	12	Shattercane	6	8
Foxtail, green	6	12	Signalgrass, broadleaf	3	5
Foxtail, robust purple	6	8	Sprangletop	4	6
Foxtail, yellow <sup>2</sup>	3	4	Sorghum, volunteer	6	8
Goosegrass <sup>3</sup>	2	3	Stinkgrass	4	6
Johnsongrass, seedling	3	5	Wheat, volunteer <sup>2</sup>	4	5
Junglerice	3	5	Witchgrass	4	6

\*Suppression only.

<sup>1</sup>Volunteer corn or rice will be best controlled with a timeline cultivation 7 - 10 days following application, and/or an additional treatment 10 - 21 days after initial application.

<sup>2</sup>Treat before tiller initiation for optimum control of crabgrass, field sandbur, wild oats, or yellow foxtail.

<sup>3</sup>For control, a sequential application may be needed.

### USE DIRECTIONS FOR POTATO VINE DESICCATION

**Axill Solutions Glufosinate 280SL** can be used to desiccate potato vines once the vines reach senescence.

#### Use Rate and Timing

Apply **Axill Solutions Glufosinate 280SL** at the rate of 21 fl. oz./A (0.38 lb. a.i./A). If a particular potato variety has a heavy or dense vine, an application of another desiccation product may be necessary for total desiccation of the potato vine.

Apply the indicated amount of **Axill Solutions Glufosinate 280SL** in enough water (20 - 100 gals. per acre) to thoroughly cover the potato vines. Take into account the density of the potato vine and increase or decrease spray volume as necessary to achieve complete coverage.

#### Important:

- It is essential to obtain thorough coverage of the potato vine for adequate desiccation. Make sure the spray boom is operated as low as possible to achieve thorough coverage (this also minimizes any potential for drift)
- If climate conditions are cool and dry, or if potato vine canopy is heavy, make sure to use a spray volume of at least 30 gals. of water per acre

**Restrictions:**

- **DO NOT** apply more than 21 fl. oz./A (0.38 lb. a.i./A) of **Axill Solutions Glufosinate 280SL** to potato vines per year.
- **DO NOT** apply more than 21 fl. oz./A (0.38 lb. a.i./A) in a single application.
- **DO NOT** apply a split application.
- **DO NOT** apply more than 1 application per year.
- Apply after the potato vine enters its natural senescence period
- Pre-harvest interval is 9 days.
- The product can be applied to potatoes grown for seed.

**Table 10** – Crop rotation and plantback intervals after application of **Axill Solutions Glufosinate 280SL** for potato vine desiccation are as follows:

Crop	Minimum Rotation Interval
All crops other than those listed in this table	120 Days
Barley, Buckwheat, Millet, Oats, Rye, Sorghum, Triticale, Wheat	30 Days
Canola, Corn, Cotton, Soybean, Sugar Beets	0 days

**USE DIRECTIONS FOR CANOLA SEED PROPAGATION**

During canola seed propagation, to eliminate vulnerable canola segregates that are not resistant to glufosinate-ammonium, apply **Axill Solutions Glufosinate 280SL** as a foliar spray, as indicated in the chart below. If canola is injured or adversely affected by environmental stress (for example: excessive rainfall, drought, pest pressure, etc.) or prior herbicide application, **DO NOT** apply **Axill Solutions Glufosinate 280SL**. If canola is injured or adversely affected by environmental stress (for example: excessive rainfall, drought, pest pressure, etc.) or prior herbicide application, **DO NOT** apply **Axill Solutions Glufosinate 280SL**.

**Table 11: RESTRICTIONS – Canola Seed Propagation**

Axill Solutions Glufosinate 280SL Use Rate		Canola Growth Stage
APPLICATION 1	22 fl. oz./A (0.4 lb. a.i./A)	Cotyledon stage up to early bolt stage (BBCH 18-30; just prior to stem elongation/bolting (8 or more leaves) through beginning of stem elongation (no internode)).
APPLICATION 2	22 fl. oz./A (0.4 lb. a.i./A)	
APPLICATION 3	22 fl. oz./A (0.4 lb. a.i./A)	

**Restrictions:**

- **DO NOT** make more than 3 applications of **Axill Solutions Glufosinate 280SL** to canola per year.
- **DO NOT** reapply within 10 days of previous application.
- **DO NOT** apply more than 22 fl. oz./A (0.4 lb. a.i./A) per application.
- **DO NOT** apply more than 66 fl. oz./A (1.2 lbs. a.i./A) per year.
- Pre-harvest interval is 65 days.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** use treated canola seed for food, feed, or oil purposes.

**USE DIRECTIONS FOR CORN, COTTON, AND SOYBEAN SEED PROPAGATION**

During seed propagation, vulnerable “segregates” (plants not resistant to glufosinate-ammonium) of corn, cotton and soybean can be selected out by application of **Axill Solutions Glufosinate 280SL**.

**CORN**

To pick out resistant segregates, apply **Axill Solutions Glufosinate 280SL** as indicated in the chart below. Corn plants not resistant to glufosinate-ammonium will be seriously injured or killed. To protect plants from **Axill Solutions Glufosinate 280SL**, apply herbicide with a hooded sprayer.

**Table 12: RESTRICTIONS – Corn Seed Propagation**

Axill Solutions Glufosinate 280SL Use Rate		Additive*	Corn Growth Stage
APPLICATION 1	22 fl. oz./A (0.4 lb. a.i./A)	AMS - 3 lbs./A	V3 - V4 (3 - 4 developed collars)
APPLICATION 2	22 fl. oz./A (0.4 lb. a.i./A)	AMS - 3 lbs./A	V6 - V7

\*AMS - Ammonium Sulfate; Reduce rate of AMS to 1.5 lbs./A when temperatures exceed 85°F, to limit possibility of leaf burn.

**Restrictions:**

- **DO NOT** make more than 2 applications per year of **Axill Solutions Glufosinate 280SL** at 22 fl. oz./A (0.4 lb. a.i./A).
- **DO NOT** apply more than 22 fl. oz./A (0.40 lb. a.i./A) in a single application.
- **DO NOT** exceed the max yearly application rate of 44 fl. oz./A (0.8 lb. a.i./A).
- **DO NOT** reapply within 10 days of previous application.

**COTTON**

During cotton seed propagation, to eliminate vulnerable cotton segregates that are not resistant to glufosinate-ammonium, apply **Axill Solutions Glufosinate 280SL** as a foliar spray, as indicated in the chart below. Two or three applications can be made, at least 10 days apart. Timing 1 can be used if weeds are particularly large or weed pressure is high due to environmental conditions preventing timely use of **Axill Solutions Glufosinate 280SL**. Use Timing 2 when making a timely application, under normal pest pressure.

**Table 13: TIMING 1 RESTRICTIONS – Cotton Seed Propagation**

Axill Solutions Glufosinate 280SL Use Rate		Cotton Growth Stage
APPLICATION 1	30 - 43 fl. oz./A (0.55 - 0.79 lb. a.i./A)	Emergence, up to early bloom.
APPLICATION 2	22 - 29 fl. oz./A (0.4 - 0.53 lb. a.i./A)	

**Restrictions:**

- **DO NOT** make more than 2 applications per year of **Axill Solutions Glufosinate 280SL**.
- **DO NOT** apply more than 43 fl. oz./A (0.34 lb. a.i./A) in a single application.
- **DO NOT** exceed the max yearly application rate of 72 fl. oz./A (1.26 lbs. a.i./A) when applied at reduced rates.
- **DO NOT** reapply within 10 days of previous application.

**Table 14: TIMING 2 RESTRICTIONS – Cotton Seed Propagation**

Axill Solutions Glufosinate 280SL Use Rate		Cotton Growth Stage
APPLICATION 1	22 - 29 fl. oz./A (0.4 - 0.53 lb. a.i./A)	Emergence, up to early bloom.
APPLICATION 2	22 - 29 fl. oz./A (0.4 - 0.53 lb. a.i./A)	
APPLICATION 3	22 - 29 fl. oz./A (0.4 - 0.53 lb. a.i./A)	

**Restrictions:**

- **DO NOT** make more than 3 applications of **Axill Solutions Glufosinate 280SL** at 29 fl. oz./A (0.53 lb. a.i./A).
- **DO NOT** apply more than 29 fl. oz./A (0.53 lb. a.i./A) in a single application.
- **DO NOT** exceed the max yearly application rate of 87 fl. oz./A (1.59 lbs. a.i./A).
- **DO NOT** reapply within 10 days of previous application.

**SOYBEANS**

To pick out resistant soybean segregates, apply **Axill Solutions Glufosinate 280SL** as indicated in the chart below. A second application, at least 5 days later, may be used, if needed.

**Table 15: RESTRICTIONS – Soybean Seed Propagation**

Axill Solutions Glufosinate 280SL Use Rate		Soybean Growth Stage
APPLICATION 1	22 fl. oz./A (0.4 lb. a.i./A)	Third trifoliolate stage.
APPLICATION 2	22 fl. oz./A (0.4 lb. a.i./A)	Up to (but not including) bloom.

**Restrictions:**

- **DO NOT** make more than 2 applications of **Axill Solutions Glufosinate 280SL** within a 1-year period.
- **DO NOT** apply more than 22 fl. oz./A (0.4 lb. a.i./A) per application.
- **DO NOT** exceed the max yearly application rate of 44 fl. oz./A (0.8 lb. a.i./A).
- **DO NOT** reapply within 10 days of previous application.

**USE DIRECTIONS FOR NON-CROP USES**

**Axill Solutions Glufosinate 280SL** is a non-selective water-soluble herbicide for application as a foliar spray for the control of a broad spectrum of emerged annual and perennial grass and broadleaf weeds. Plants that have not yet emerged at the time of application will not be controlled. THOROUGH SPRAY COVERAGE IS IMPORTANT. Visual effects and control from application of this product occur within 2 - 4 days after application under good growing conditions.

This product is non-selective and will injure or kill all green vegetation contacted by the spray. **DO NOT** allow contact with foliage or green tissue of desirable vegetation. **DO NOT** allow direct spray or drift onto green, thin, or uncalloused bark of desirable vegetation or plant injury may result. If desirable vegetation is contacted, rinse sprayed portion with water immediately.

**WHERE TO APPLY**

**Axill Solutions Glufosinate 280SL** may be used on the following: access roads, airfields, airports, alleys, along fences and fence lines, around commercial or industrial structures or outbuildings, around farm and ranch structures and outbuildings, around ornamental gardens, around ornamental trees and shrubs (including Christmas trees), bare ground, barrier strips, beaches\*, campgrounds, canals, commercial/industrial plant sites, Conservation Reserve Program (CRP)\*, construction sites, ditches and ditch banks, drive-in theaters, driveways, dry ditches, fencerows, firebreaks, fuel storage areas, golf courses (excluding greens, tees, aprons, fairways, and roughs)\*, gravel yards, habitat restoration and management areas, highways and roadsides (including aprons, medians, guardrails, and right of ways), industrial areas, industrial plant sites, landscapes, lanes, lumberyards, mulched areas, natural areas and brush control, nurseries, parking areas, parks, paths, paved areas, petroleum and other tank farms, pipeline, power, telephone, and utility rights of way, power stations, pre-plant to turf and ornamental plants, pumping installations, pumping stations, railroad rights of way, ramps, recreation areas, refineries, sewage disposal areas, shelter belts, sidewalks, site preparation areas for conifer and hardwood, storage areas, substations, tennis courts, trails, uncropped farmstead areas, vacant lots, walkways, wastelands, wildlife habitat areas, wildlife openings.

\*Not for use in California.

### Restrictions For Non-Crop Use:

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation.
- **DO NOT** allow grazing of vegetation treated with this product.
- **DO NOT** apply more than 246 fl. oz. of this product per acre per year (4.5 lbs. a.i./A/year).
- **DO NOT** apply more than 82 fl. oz. of this product per acre per single application (1.5 lbs. a.i./A/application).
- **DO NOT** apply more than a total of 3 broadcast applications per year.
- **DO NOT** apply more than 2 applications per year on Dormant bermudagrass.
- **DO NOT** reapply within 14 days of previous application in non-crop areas.
- **DO NOT** apply to residential areas or areas where crops may be grown.

### Restrictions for Non-Crop Use - Spot or Directed Applications:

- **DO NOT** apply beyond runoff.
- **DO NOT** spray during windy conditions.
- **DO NOT** exceed single maximum and yearly maximum broadcast use rates.
- **DO NOT** make more than 3 spot treatment applications to same 1,000 sq. ft. area per year.
- **DO NOT** apply more than 82 fl. oz. of product per acre (1.5 lbs a.i./A) or 1.9 fl. oz. of product per 1,000 sq. ft. (0.03 lbs a.i./1,000 sq. ft.) per single spot treatment.
- **DO NOT** apply more than 246 fl. oz. of this product per acre (4.5 lbs. a.i./A) or 5.7 fl. oz. of product per 1,000 sq. ft. (0.09 lbs a.i.) per year.

### WHEN TO APPLY

**Axill Solutions Glufosinate 280SL** is a foliar-active material and works best when weeds are actively growing. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application of the highest rate indicated.

Always apply this product at the labeled rate in **HOW TO APPLY**. Repeat applications or tank mixes of **Axill Solutions Glufosinate 280SL** plus 1 or more appropriate residual herbicides will be needed to control weeds emerging from underground parts or seeds.

### MIXING INSTRUCTIONS

**Axill Solutions Glufosinate 280SL** must be mixed with water to make a finished spray solution. Fill the spray tank with water, start agitation, add the appropriate amount of product then mix thoroughly.

### HOW TO APPLY

#### Spot or Directed Applications

Apply 0.4 - 0.75 fl. oz. (0.0007 - 0.014 lb. a.i.) per gallon of water. Use rate depends on weed and stage of growth as shown in the following sections. Spray undesirable vegetation foliage on a spray-to-wet basis. Ensure uniform and complete coverage. Use a coarse spray. Backpack, pump-up, and hydraulic sprayers may be used. Thoroughly clean the sprayer following use.

#### Restrictions:

#### Broadcast or Boom Applications

Apply 12 - 38 fl. oz. (0.22 - 0.69 lb. a.i./A) per acre in a minimum of 40 gals. of water. Use rate depends on weed and stage of growth in the following section. Use 30-PSI spray pressure minimum.

#### Aerial Applications

Apply as a foliar treatment using a minimum of 5 gals. of water per acre to ensure thorough coverage. **DO NOT** apply when winds are gusty or under conditions which favor drift on to desirable vegetation. Applications under conditions which cause drift of this product will result in damage to any vegetation contacted. Drift control additives maybe used. If a drift control additive is used, observe, and follow all directions and precautions as specified on the additive label.

### Tank Mix Directions for Non-Crop Uses

**Axill Solutions Glufosinate 280SL** is compatible in tank mixes with many other herbicides including non-selective herbicides. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Tank mix applications of **Axill Solutions Glufosinate 280SL** plus the following herbicides are advised for broad-spectrum post-emergence and pre-emergence weed control:

Isopropylamine salt of imazapyr	butroxydim	norflurazon
prodiamine	isoxaben	Diglycolamine salt of 3,6-dichloro-o-anisic Acid
oryzalin	pendimethalin	oxadiazon

A compatibility test must be conducted with any potential tank mix partner with **Axill Solutions Glufosinate 280SL**, except with any 1 of those listed above. Using a clear glass quart jar, conduct the test as described below:

1. Fill the jar 3/4 full with water.
2. Add the appropriate amount of herbicide in the following order: (a) dry flowable, (b) wettable powder, (c) aqueous suspensions, (d) flowables, (e) liquids and (f) solutions and emulsifiable or liquid concentrates. Shake or gently stir jar after each addition to thoroughly mix.

3. After adding all ingredients, let the mixture stand for 15 minutes and then look for separation, large flakes, precipitates, gels, and heavy oily film on the jar or other signs of incompatibility.
4. If the compatibility test shows signs of incompatibility, tank mix the product tested with **Axill Solutions Glufosinate 280SL**.

**For the following Weeds Controlled by Axill Solutions Glufosinate 280SL Apply:**

- **Broadleaf Weeds:** Chickweed, Clover, Common cocklebur, Filaree, Jimsonweed, Kochia, London rocket, Malva (little mallow), Maretail, Purslane, Shepherd's purse, Smartweed
- **Grasses and Sedges:** Barnyardgrass, Cupgrass, Fall Panicum, Giant Foxtail, Goosegrass, Green Foxtail, Johnsongrass (rhizome), Lovegrass, Shattercane, Smallflower Alexandergrass (Signalgrass), stinkgrass, Windgrass, Yellow foxtail
- **Spot Application (1,000 sq. ft.):** Apply 0.75 fl. oz./gal. of water (0.014 lb. a.i./gal. of water) when the weed height or diameter is less than 6 inches. Apply 1.25 fl. oz./ gallon of water (0.023 lb. a.i./gal. of water) when the weed height or diameter is 6 inches or greater.
- **Broadcast Application:** Apply 40 fl. oz./A (0.73 lb. a.i./A) when the weed height or diameter is less than 6 inches. Apply 56 fl. oz./A (1.02 lbs. a.i./A) when the weed height or diameter is 6 inches or greater.

**For The Following Weeds Controlled by Axill Solutions Glufosinate 280SL Apply:**

- **Broadleaf Weeds:** Annual sowthistle, Bindweed, Buffalobur, Burdock, Canada thistle, Curly dock, Dandelion, Dogbane, (hemp), Field gromwell, Fleabane, Goldenrod, Horsetail, Lambsquarters, Leafy spurge, Mugwort, Musk thistle, Nettle, Nightshade, Pennycress, Pigweed (redroot), Plantain, Prickly lettuce, ragweed, Russian thistle, Tansy mustard, Velvetleaf, Verbain, Virginia copperleaf, White heath aster, Wild buckwheat, Wild mustard, Wild onion, Wild rose, Wild turnip, Woodsorrel, Yellow rocket
- **Grasses and Sedges:** Annual bluegrass, Bahiagrass, Barley, Bermudagrass, Carpetgrass, Crabgrass, Dallisgrass, Downy bromegrass, Fescue, Guineagrass, Kentucky bluegrass, nutsedge, Paragrass, Quackgrass, Ryegrass, Sandbur, Smooth Bromegrass, Torpedograss, Vaseygrass, Wheat, Wild oat
- **Spot Application (1,000 sq. ft.):** Apply 1.25 fl. oz./gal. of water (0.023 lb. a.i./gal. of water) when the weed height or diameter is less than 6 inches. Apply 1.75 fl. oz./gal. of water (0.032 lb. a.i./gal. of water) when the weed height or diameter is 6 inches or greater.
- **Broadcast Application:** Apply 56 fl. oz./A (1.02 lbs. a.i./A) when the weed height or diameter is less than 6 inches tall. Apply 80 fl. oz./A (1.46 lbs. a.i./A) when the weed height or diameter is 6 inches or greater.

**Additional Use Directions**

1. Use higher rates within the directed rate range for plant sizes listed when vegetation cover is dense or when weeds are grown under stressed conditions including drought or when average temperature are below 50°F.
2. The addition of 8.5 - 17 lbs. of ammonium sulfate (spray grade) per 100 gals. of water (1% - 2% by weight) or 2 - 4 lbs. of ammonium sulfate per acre may improve the level of weed control.

**Use on Woody Species**

**(Not for use in California)**

When applied as labeled, **Axill Solutions Glufosinate 280SL** will provide control, partial control, or suppression of certain perennial woody weed species. Apply 64 - 82 fl. oz./A (1.19 - 1.5 lbs. a.i./A). Use the higher specified rates per acre of this product when conditions are not optimum for spray penetration, including when vegetation growth is heavy or dense. Lower specified rates may be used when the target species is a conifer and when vegetation growth conditions allow for uniform spray coverage.

Blackberry	<i>Rubus</i> spp.
Deer brush	<i>Ceanothus integerrimus</i>
Douglas fir	<i>Pseudotsuga menziesii</i>
Gallberry	<i>Ilex</i> spp.
Hazel	<i>Corylus</i> spp.
Honeysuckle	<i>Lonicera</i> spp.
Huckleberry	<i>Gaylussacia</i> spp.
Maple	<i>Acer</i> spp.
Multiflora rose	<i>Rosa multiflora</i>
Oak	<i>Quercus</i> spp.
Pine	<i>Pinus</i> spp.
Poison ivy	<i>Toxicodendron radicans</i>
Poison oak	<i>Toxicodendron toxicarium</i>
Roundleaf greenbrier	<i>Smilax rotundifolia</i>
Salmonberry	<i>Rubus spectabilis</i>
Sweet gum	<i>Liquidambar styraciflua</i>
Sumac	<i>Rhus</i> spp.
Thimbleberry	<i>Rubus parviflorus</i>
Trumpet creeper	<i>Campsis radicans</i>
Vine maple	<i>Acer circinatum</i>
Western red cedar	<i>Thuja plicata</i>

### **Conservation Reserve Program (CRP)\***

**Axill Solutions Glufosinate 280SL** may be used to suppress competitive growth and seed production of undesirable vegetation when rotating out of CRP acres. Apply 48 - 56 fl. oz. (0.88 - 1 lb. a.i.) per acre of **Axill Solutions Glufosinate 280SL** in early spring, before CRP grasses break dormancy and initiate growth. Late fall applications may be made after desirable perennial grasses have reached dormancy. Some stunting of CRP perennial grasses may occur if applications are made when plants are not dormant.

\*Not for use in California

### **Trimming and Edging**

**Axill Solutions Glufosinate 280SL** may be used for trimming and edging areas listed under the **WHERE TO APPLY** section. For control of weeds emerging from seed, the use of **Axill Solutions Glufosinate 280SL** in a tank mix with pre-emergence herbicides is advised. If spraying in areas adjacent to desirable plants, use a shield made of cardboard, plywood, or sheet metal while spraying to help prevent spray from contacting foliage of desirable plants. Refer to the **HOW TO APPLY** section below for appropriate application rates to control specific weeds.

### **Wildlife Food Plots\***

This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seed bed, wait 7 days after applying this product before tilling. Refer to the **HOW TO APPLY** section of this labeling for appropriate application rates to control specific weeds.

\*Not for use in California

### **Dormant Bermudagrass (Not for use on Residential Turf/Turfgrass/Lawns)**

**Axill Solutions Glufosinate 280SL** may be used to control winter annual weeds in well-established ornamental dormant hybrid or common Bermudagrass. Apply only when the turf is fully dormant and prior to spring green-up or severe turfgrass injury or delayed green-up may occur. For best results, apply this product at a rate of 40 - 80 fl. oz. (0.73 - 1.46 lbs. a.i.) per acre after most weeds have germinated and are in an early growth stage. Referred to the **HOW TO APPLY** section below for specified rates. Applications of this product may also be used to suppress or control target biennial or perennial weeds. Avoid high volume and spot applications where spray volume exceeds 80 gals. per acre or injury or delayed green-up may occur.

### **Restrictions:**

- **DO NOT** apply more than 80 fl. oz. (1.46 lbs. a.i.)/A per year for this use.
- **DO NOT** use on Residential Turf/Turfgrass/Lawns

### **Ornamentals and Christmas Trees**

When applied as advised by this label, this product may be used for the control of undesired vegetation in site preparation prior to planting, around and within shade and greenhouses, and as a directed spray around containers and field-grown established ornamentals and Christmas Trees. Refer to the **HOW TO APPLY** section below for appropriate application rates to control specific weeds.

- **Pre-Plant Site Preparation:** **Axill Solutions Glufosinate 280SL** may be used for pre-plant site preparation for the control of annual and perennial weeds listed on this label in ornamental and Christmas Tree plantings. Refer to the **HOW TO APPLY** section below for appropriate application rates to control specific weeds.
- **Directed Spray Application:** **Axill Solutions Glufosinate 280SL** may be applied as a directed spray to control in-row weeds in field-grown woody plants. Refer to the **HOW TO APPLY** section below for appropriate application rate to control specific weeds. This product may be used between and around container and site preparation for new planting.

### **Restrictions:**

- Plant Ornamental and Christmas Trees only after the restricted entry interval (REI) of 12 hours.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation or injury will result.
- **DO NOT** apply this product as an over-the-top broadcast spray in ornamentals and shade or Christmas trees.

### **Greenhouse and Shade House Applications**

**Axill Solutions Glufosinate 280SL** may be used to control weeds in greenhouses and shade-houses. Air circulation fans must be turned off during application. Apply this product as a directed spray using large droplet and low-pressure type nozzles. Avoid drift and direct contact with desirable vegetation.

**Restriction:** **DO NOT** use in greenhouses or shade houses containing edible crops.

### **Site Preparation for Conifer and Hardwood Production Areas**

Prior to planting conifer and hardwood species, **Axill Solutions Glufosinate 280SL** can be used as a site preparation treatment.

### **Restrictions:**

- **DO NOT** apply **Axill Solutions Glufosinate 280SL** as an over-the-top broadcast spray to desirable conifer or hardwood plantings.
- Restricted Entry Interval (REI) for seedling conifer and hardwood trees to be planted into the treated area is 12 hours.

# STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE: DO NOT** use or store near heat or open flame. Keep the container tightly closed and dry in a cool, well-ventilated place. **DO NOT** exceed storage temperature of 125°F. If storage temperature for bulk product is below 32°F, **DO NOT** pump material until its temperature exceeds 32°F. Protect against direct sunlight.

**PESTICIDE DISPOSAL:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

## CONTAINER HANDLING:

**Less Than or Equal to 5 Gallons:** Non-refillable plastic container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

**Greater Than 5 Gallons:** Non-refillable plastic container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

## CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Axill Solutions, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Axill Solutions, LLC and Seller harmless for any claims relating to such factors.

Axill Solutions, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Axill Solutions, LLC, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, AXILL SOLUTIONS, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Axill Solutions, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF AXILL SOLUTIONS, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF AXILL SOLUTIONS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Axill Solutions, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Axill Solutions, LLC.